

Table des matières

1. Systematic Reviews and Meta-Analysis	1
1.1. Acupuncture générique	1
1.1.1. Liao 2021	1
1.1.2. Li 2009 ☆☆	2
1.1.3. Liu 2009	2
1.1.4. Van Den Berg 2008 ☆☆	3
1.1.5. Van den Berg 2005	3
1.2. Specific Acupuncture Techniques	3
1.2.1. Moxibustion	3
1.2.1.1. Coyle 2023	3
1.2.1.2. Zhang 2019 (æ67V)	6
1.2.1.3. Wu 2018	6
1.2.1.4. Zhang 2013 ☆☆	6
1.2.1.5. Coyle 2012 ☆	7
1.2.1.6. Vas 2009 ☆☆	8
1.2.1.7. Coyle 2005 Ø	8
2. Cost-effectiveness	9
3. Clinical Practice Guidelines	9
3.1. National Institute for Health and Care Excellence (NICE, UK) 2021 Ø	9
3.2. Danish Society of Obstetrics and Gynecology (DSOG, Denmark) 2020 ⊕	9
3.3. Collège national des gynécologues et obstétriciens français (CNGOF, France) 2019 Ø	10
3.4. Royal College of Obstetricians and Gynaecologists (RCOG, UK) 2017 ⊕	10
4. Overviews of Clinical Practice Guidelines	10
4.1. Tsakiridis 2019	10
5. Randomized Controlled Trials	11
5.1. Sources	11
5.2. List	11

breech presentation

Version foetale : évaluation de l'acupuncture

1. Systematic Reviews and Meta-Analysis

GRADE ratings and their interpretation

Symbol	Quality	Interpretation
⊕⊕⊕⊕	High	We are very confident that the true effect lies close to that of the estimate of the effect.
⊕⊕⊕⊖	Moderate	We are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.
⊕⊕⊖⊖	Low	Our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.
⊕⊖⊖⊖	Very low	We have very little confidence in the effect estimate: the true effect is likely to be

1.1. Acupuncture générique

1.1.1. Liao 2021

Liao JA, Shao SC, Chang CT, Chai PYC, Owang KL, Huang TH, Yang CH, Lee TJ, Chen YC. Correction of Breech Presentation with Moxibustion and Acupuncture: A Systematic Review and Meta-Analysis. *Healthcare*. 2021;9(6):619. [212237]. [doi](#)

Aim	Acupuncture-type interventions (such as moxibustion and acupuncture) at Bladder 67 (BL67, Zhiyin point) have been proposed to have positive effects on breech presentation. The aim of this systematic review and meta-analysis was to evaluate the effectiveness and safety of moxibustion and acupuncture in correcting breech presentation.
Methods	We searched PubMed, MEDLINE, Embase, the Cochrane Central Register of Controlled Trials (CENTRAL), the Chinese Electronic Periodical Services (CEPS), and databases at ClinicalTrials.gov to identify relevant randomized controlled trials (RCTs).
Results	In this study, sixteen RCTs involving 2555 participants were included. Compared to control, moxibustion significantly increased cephalic presentation at birth (RR = 1.39; 95% CI = 1.21-1.58). Moxibustion also seemed to elicit better clinical outcomes in the Asian population (RR = 1.42; 95% CI = 1.21-1.67) than in the non-Asian population (RR = 1.20; 95% CI = 1.01-1.43). The effects of acupuncture on correcting breech presentation after sensitivity analysis were inconsistent relative to control. The effect of moxibustion plus acupuncture was synergistic for correcting breech presentation (RR = 1.53; 95% CI = 1.26-1.86) in one RCT.
Conclusions	Our findings suggest that moxibustion therapy has positive effects on correcting breech presentation, especially in the Asian population.

1.1.2. Li 2009 ☆☆

Li X, Hu J, Wang X, Zhang H, Liu J. Moxibustion and other acupuncture point stimulation methods to treat breech presentation: a systematic review of clinical trials. *Chin Med.* 2009;4(1):4.[153149].

Background	Moxibustion, acupuncture and other acupoint stimulations are commonly used for the correction of breech presentation. This systematic review aims to evaluate the efficacy and safety of moxibustion and other acupoint stimulations to treat breech presentation.
Methods	We included randomized controlled trials (RCTs) and controlled clinical trials (CCTs) on moxibustion, acupuncture or any other acupoint stimulating methods for breech presentation in pregnant women. All searches in PubMed, the Cochrane Library (2008 Issue 2), China National Knowledge Information (CNKI), Chinese Scientific Journal Database (VIP) and WanFang Database ended in July 2008. Two authors extracted and analyzed the data independently.
Results	Ten RCTs involving 2090 participants and seven CCTs involving 1409 participants were included in the present study. Meta-analysis showed significant differences between moxibustion and no treatment (RR 1.35, 95% CI 1.20 to 1.51; 3 RCTs). Comparison between moxibustion and knee-chest position did not show significant differences (RR 1.30, 95% CI 0.95 to 1.79; 3 RCTs). Moxibustion plus other therapeutic methods showed significant beneficial effects (RR 1.36, 95% CI 1.21 to 1.54; 2 RCTs). Laser stimulation was more effective than assuming the knee-chest position plus pelvis rotating. Moxibustion was more effective than no treatment (RR 1.29, 95% CI 1.17 to 1.42; 2 CCTs) but was not more effective than the knee-chest position treatment (RR 1.22, 95% CI 1.11 to 1.34; 2 CCTs). Laser stimulation at Zhiyin (BL67) was more effective than the knee-chest position treatment (RR 1.30, 95% CI 1.10 to 1.54; 2 CCTs,).
Conclusion	Moxibustion, acupuncture and laser acupoint stimulation tend to be effective in the

1.1.3. Liu 2009

Liu Mai-Lan, Lan Lei, Tang Yong, Liang Fan-Rong. [Acupuncture and moxibustion for breech presentation: a systematic review]. *Chinese Journal of EBM.* 2009;9(8):840-3. [167341].

Objective	To evaluate the efficacy of acupuncture and moxibustion (acup-moxi) therapy for breech presentation.
Methods	We electronically searched The Cochrane Library (Issue 1, 2008), PubMed (1980 to Mar. 2008), MEDLINE (1966 to 2008), Ovid EBM Database (1991 to 2008), CBMdisc (1978 to Mar. 2008), VIP (1989 to Mar. 2008), CNKI (1979 to Mar. 2008), and WangFang Database (1983 to Mar. 2008), as well as handsearched seven traditional Chinese medicine journals to obtain randomized control trials (RCTs) about acup-moxi for breech presentation. Quality assessment was conducted according to the Cochrane Handbook for Systematic Reviews of Interventions 5.0.1. Metaanalyses were performed for the results of homogeneous studies using RevMan 5.0 software.
Results	Eight RCTs involving 1 341 patients met the inclusion criteria. Five trials were of relatively high quality and 3 were of low quality. The pooled analysis of six trials showed that acup-moxi was superior in cephalic presentation with RR=1.38, and 95%CI 1.20 to 1.58.
Conclusion	Acup-moxi can increase the successful rate of cephalic presentation in the treatment of breech presentation compared with no (routine care) or knee-chest position treatment.

1.1.4. Van Den Berg 2008 ☆☆

Van Den Berg I et al. Effectiveness of acupuncture-type interventions versus expectant management to correct breech presentation: a systematic review. Complement Ther Med. 2008;16(2):92-100.149428

Purpose	A systematic review of studies assessing the effectiveness of acupuncture-type interventions (moxibustion, acupuncture, or electro-acupuncture) on acupuncture point BL 67 to correct breech presentation compared to expectant management, based on controlled trials.
Methods	Articles published from 1980 to May 2007 in databases of Medline, EMBASE, the Cochrane Central Register of Controlled Trials, AMED, NCCAM, Midirs and reference lists. Study selection: studies included were original articles; randomised controlled trials (RCT or controlled cohort studies; acupuncture-type intervention on BL 67 compared with expectant management; ultrasound confirmed breech presentation and position of the fetus after treatment confirmed with ultrasound, position at delivery, and/or the proportion of caesarea sections reported.
Results	Of 65 retrieved citations, six RCT's and three cohort studies fulfilled the inclusion criteria. Data were pooled using random-effects models. In the RCT's the pooled proportion of breech presentations was 34% (95% CI: 20-49%) following treatment versus 66% (95% CI: 55-77%) in the control group (OR 0.25 95% CI: 0.11-0.58). The pooled proportion in the cohort studies was 15% (95% CI: 1-28%) versus 36% (95% CI: 14-58%), (OR 0.29, 95% CI: 0.19-0.43). Including all studies the pooled proportion was 28% (95% CI: 16-40%) versus 56% (95% CI: 43-70%) (OR 0.27, 95% CI: 0.15-0.46).
Conclusion	Our results suggest that acupuncture-type interventions on BL 67 are effective in correcting breech presentation compared to expectant management. Some studies were of inferior quality to others and further RCT's of improved quality are necessary to adequately answer the research question.

1.1.5. Van den Berg 2005

Van Den Berg I, Jacobs B, Bouman I, Bosch JL, Hunink MGM. Effectiveness of acupuncture-type interventions to resolve breech presentation compared to expectative policy, a meta-analysis. Focus on Alternative and Complementary Therapies. 2005;10(1):9. [144431].

1.2. Specific Acupuncture Techniques

1.2.1. Moxibustion

1.2.1.1. Coyle 2023

⊕⊕⊕⊖	Moderate
⊕⊕⊖⊖	Low

Coyle ME, Smith C, Peat B. Cephalic version by moxibustion for breech presentation. Cochrane Database Syst Rev. 2023 May 9;5:CD003928. <https://doi.org/10.1002/14651858.cd003928.pub4>

Background	Breech presentation at term can cause complications during birth and increase the chance of caesarean section. Moxibustion (a type of Chinese medicine which involves burning a herb close to the skin) at the acupuncture point Bladder 67 (BL67) (Chinese name Zhiyin), located at the tip of the fifth toe, has been proposed as a way of changing breech presentation to cephalic presentation. This is an update of a review first published in 2005 and last published in 2012.
Objectives	To examine the effectiveness and safety of moxibustion on changing the presentation of an unborn baby in the breech position, the need for external cephalic version (ECV), mode of birth, and perinatal morbidity and mortality.
Methods	Search methods: For this update, we searched Cochrane Pregnancy and Childbirth's Trials Register (which includes trials from CENTRAL, MEDLINE, Embase, CINAHL, and conference proceedings), ClinicalTrials.gov, and the WHO International Clinical Trials Registry Platform (ICTRP) (4 November 2021). We also searched MEDLINE, CINAHL, AMED, Embase and MIDIRS (inception to 3 November 2021), and the reference lists of retrieved studies. Selection criteria: The inclusion criteria were published and unpublished randomised or quasi-randomised controlled trials comparing moxibustion either alone or in combination with other techniques (e.g. acupuncture or postural techniques) with a control group (no moxibustion) or other methods (e.g. acupuncture, postural techniques) in women with a singleton breech presentation. Data collection and analysis: Two review authors independently determined trial eligibility, assessed trial quality, and extracted data. Outcome measures were baby's presentation at birth, need for ECV, mode of birth, perinatal morbidity and mortality, maternal complications and maternal satisfaction, and adverse events. We assessed the certainty of the evidence using the GRADE approach.

<p>Main Results</p>	<p>This updated review includes 13 studies (2181 women), of which six trials are new. Most studies used adequate methods for random sequence generation and allocation concealment. Blinding of participants and personnel is challenging with a manual therapy intervention; however, the use of objective outcomes meant that the lack of blinding was unlikely to affect the results. Most studies reported little or no loss to follow-up, and few trial protocols were available. One study that was terminated early was judged as high risk for other sources of bias. Meta-analysis showed that compared to usual care alone, the combination of moxibustion plus usual care probably reduces the chance of non-cephalic presentation at birth (7 trials, 1152 women; risk ratio (RR) 0.87, 95% confidence interval (CI) 0.78 to 0.99, I² = 38%; moderate-certainty evidence), but the evidence is very uncertain about the effect of moxibustion plus usual care on the need for ECV (4 trials, 692 women; RR 0.62, 95% CI 0.32 to 1.21, I² = 78%; low-certainty evidence) because the CIs included both appreciable benefit and moderate harm. Adding moxibustion to usual care probably has little to no effect on the chance of caesarean section (6 trials, 1030 women; RR 0.94, 95% CI 0.83 to 1.05, I² = 0%; moderate-certainty evidence). The evidence is very uncertain about the effect of moxibustion plus usual care on the the chance of premature rupture of membranes (3 trials, 402 women; RR 1.31, 95% CI 0.17 to 10.21, I² = 59%; low-certainty evidence) because there were very few data. Moxibustion plus usual care probably reduces the use of oxytocin (1 trial, 260 women; RR 0.28, 95% CI 0.13 to 0.60; moderate-certainty evidence). The evidence is very uncertain about the chance of cord blood pH less than 7.1 (1 trial, 212 women; RR 3.00, 95% CI 0.32 to 28.38; low-certainty evidence) because there were very few data. We are very uncertain whether the combination of moxibustion plus usual care increases the chance of adverse events (including nausea, unpleasant odour, abdominal pain and uterine contractions; intervention: 27/65, control: 0/57), as only one study presented data in a way that could be reanalysed (122 women; RR 48.33, 95% CI 3.01 to 774.86; very low-certainty evidence). When moxibustion plus usual care was compared with sham moxibustion plus usual care, we found that moxibustion probably reduces the chance of non-cephalic presentation at birth (1 trial, 272 women; RR 0.74, 95% CI 0.58 to 0.95; moderate-certainty evidence) and probably results in little to no effect on the rate of caesarean section (1 trial, 272 women; RR 0.84, 95% CI 0.68 to 1.04; moderate-certainty evidence). No study that compared moxibustion plus usual care with sham moxibustion plus usual care reported on the clinically important outcomes of need for ECV, premature rupture of membranes, use of oxytocin, and cord blood pH less than 7.1, and one trial that reported adverse events reported data for the whole sample. When moxibustion was combined with acupuncture and usual care, there was very little evidence about the effect of the combination on non-cephalic presentation at birth (1 trial, 226 women; RR 0.73, 95% CI 0.57 to 0.94) and at the end of treatment (2 trials, 254 women; RR 0.73, 95% CI 0.57 to 0.93), and on the need for ECV (1 trial, 14 women; RR 0.45, 95% CI 0.07 to 3.01). There was very little evidence about whether moxibustion plus acupuncture plus usual care reduced the chance of caesarean section (2 trials, 240 women; RR 0.80, 95% CI 0.65 to 0.99) or pre-eclampsia (1 trial, 14 women; RR 5.00, 95% CI 0.24 to 104.15). The certainty of the evidence for this comparison was not assessed.</p>
<p>Authors' conclusions</p>	<p>We found moderate-certainty evidence that moxibustion plus usual care probably reduces the chance of non-cephalic presentation at birth, but uncertain evidence about the need for ECV. Moderate-certainty evidence from one study shows that moxibustion plus usual care probably reduces the use of oxytocin before or during labour. However, moxibustion plus usual care probably results in little to no difference in the rate of caesarean section, and we are uncertain about its effects on the chance of premature rupture of membranes and cord blood pH less than 7.1. Adverse events were inadequately reported in most trials.</p>

1.2.1.2. Zhang 2019 (æ67V)

Zhang Lu, Sun Weipeng, Cai Yinhe, Lu Ke, Yang Yaqin, Zeng Lei, Chen Baoyan. [Meta-analysis of Efficacy of Moxibustion at Zhiyin Point for Breech Presentation]. Journal of Hubei University of Chinese Medicine. 2019;5:117-21. [100163].

Objective	To systematically evaluate the clinical efficacy of moxibustion at Zhiyin point alone or in combination with knee-chest decubitus position in treatment of fetal malposition.
Methods	PubMed, Cochrane library, CNKI, VIP, Wanfang and CBM were searched electronically. The search time was limited to June 15,2017. All relevant literatures about moxibustion at Zhiyin point or treatment of fetal malposition with knee-chest decubitus position were detected. Two researchers independently and strictly carried out quality evaluation and data extraction for the study. Meta-analysis was carried out with Stata 14 software, and sequential analysis was carried out with TSA v0.9 software.
Results	Totally 2331 patients were included in 12 eligible literatures. Meta-analysis showed that compared with knee-chest decubitus position, the positive rate of fetal position in simple group increased by 44%[RR=1.44,95%CI(1.33,1.55),P<0.001],while that in combined group increased by 35% [RR=1.32,95%CI(1.22,1.42),P<0.001],all with statistical significance. Subgroup analysis showed that the positive rate of fetal position in simple group or combined group was related to the course of treatment and moxibustion time,and both groups had the best curative effect on pregnant women with 30-34 gestational weeks. Sequential analysis of the tests indicated that moxibustion at Zhiyin point or combined with knee-chest decubitus was effective.
Conclusion	Moxibustion at Zhiyin point alone or combined with knee-chest decubitus position can increase the rate of positive fetal position, and moxibustion at Zhiyin point alone has better effect on correcting abnormal fetal position. Both groups have the best effect on pregnant women with 30-34 gestational weeks, which is worthy of clinical promotion

1.2.1.3. Wu 2018

Wu Yan. [A Meta-analysis of treating malpresentation by moxibustion]. Clinical Journal of Chinese Medicine. 2018;(2). [27280].

Objective	To evaluate the clinical efficacy of different moxibustion on malposition.
Methods	According to the Cochrane evaluation method, the literature of treating malposition by moxibustion were comprehensively collected; a Meta-analysis was performed statistically by Rev Man 5. 2.
Result	8 literatures were included in the study.
Conclusion	The effect of moxibustion and moxibustion plus other therapies on malposition was better than knee-chest position; moxibustion on malposition is superior to moxibustion plus postural therapy

1.2.1.4. Zhang 2013 ☆☆

Zhang QH, Yue JH, Liu M, Sun ZR, Sun Q, Han C et al. Moxibustion for the correction of nonvertex presentation: a systematic review and meta-analysis of randomized controlled trials. Evid Based Complement Alternat Med 2013. [160352].

Objectives	This study aims to assess the effectiveness and safety of moxibustion for the correction of nonvertex presentation.
-------------------	---

Methods	Records without language restrictions were searched up to February 2013 for randomized controlled trials (RCTs) comparing moxibustion with other therapies in women with a singleton nonvertex presentation. Cochrane risk of bias criteria were used to assess the methodological quality of the trials.
Results	Seven of 392 potentially relevant studies met the inclusion criteria. When moxibustion was compared with other interventions, a meta-analysis revealed a significant difference in favor of moxibustion on the correction of nonvertex presentation at delivery (risk ratio RR 1.29, 95% confidence interval (CI) 1.12 to 1.49, and I ² = 0). The same findings applied to the cephalic presentation after cessation of treatment (RR 1.36, 95% CI 1.08 to 1.71, and I ² = 80%). A subgroup analysis that excluded two trials with a high risk of bias also indicated favorable effects (RR 1.63, 95% CI 1.42 to 1.86, and I ² = 0%). With respect to safety, moxibustion resulted in decreased use of oxytocin.
Conclusion	Our systematic review and meta-analysis suggested that moxibustion may be an effective treatment for the correction of nonvertex presentation. Moreover, moxibustion might reduce the need for oxytocin.

1.2.1.5. Coyle 2012 ☆

Coyle Me, Smith Ca, Peat B. Cephalic version by moxibustion for breech presentation. Cochrane Database Syst Rev. 2012. May 16:. [159289].

Background	Moxibustion (a type of Chinese medicine which involves burning a herb close to the skin) to the acupuncture point Bladder 67 (BL67) (Chinese name Zhiyin), located at the tip of the fifth toe, has been proposed as a way of correcting breech presentation.
Objectives	To examine the effectiveness and safety of moxibustion on changing the presentation of an unborn baby in the breech position, the need for external cephalic version (ECV), mode of birth, and perinatal morbidity and mortality for breech presentation.
Methods	Search methods: We searched the Cochrane Pregnancy and Childbirth Group's Trials Register (26 March 2012), MEDLINE (1966 to 1 August 2011), EMBASE (1980 to August 2011), CINAHL (1982 to 1 August 2011), MIDIRS (1982 to 1 August 2011) and AMED (1985 to 1 August 2011) and searched bibliographies of relevant papers. Selection criteria: The inclusion criteria were published and unpublished randomised controlled trials comparing moxibustion (either alone or in combination with acupuncture or postural techniques) with a control group (no moxibustion), or other methods (e.g. external cephalic version, acupuncture, postural techniques) in women with a singleton breech presentation. DATA collection and analysis: Two review authors independently assessed eligibility and trial quality and extracted data. The outcome measures were baby's presentation at birth, need for external cephalic version, mode of birth, perinatal morbidity and mortality, maternal complications and maternal satisfaction, and adverse events.

<p>Main results</p>	<p>Six new trials have been added to this updated review. One trial has been moved to studies awaiting classification while further data are being requested. This updated review now includes a total of eight trials (involving 1346 women). Meta-analyses were undertaken (where possible) for the main and secondary outcomes. Moxibustion was not found to reduce the number of non-cephalic presentations at birth compared with no treatment (P = 0.45). Moxibustion resulted in decreased use of oxytocin before or during labour for women who had vaginal deliveries compared with no treatment (risk ratio (RR) 0.28, 95% confidence interval (CI) 0.13 to 0.60). Moxibustion was found to result in fewer non-cephalic presentations at birth compared with acupuncture (RR 0.25, 95% CI 0.09 to 0.72). When combined with acupuncture, moxibustion resulted in fewer non-cephalic presentations at birth (RR 0.73, 95% CI 0.57 to 0.94), and fewer births by caesarean section (RR 0.79, 95% CI 0.64 to 0.98) compared with no treatment. When combined with a postural technique, moxibustion was found to result in fewer non-cephalic presentations at birth compared with the postural technique alone (RR 0.26, 95% CI 0.12 to 0.56).</p>
<p>Authors' conclusions</p>	<p>This review found limited evidence to support the use of moxibustion for correcting breech presentation. There is some evidence to suggest that the use of moxibustion may reduce the need for oxytocin. When combined with acupuncture, moxibustion may result in fewer births by caesarean section; and when combined with postural management techniques may reduce the number of non-cephalic presentations at birth, however, there is a need for well-designed randomised controlled trials to evaluate moxibustion for breech presentation which report on clinically relevant outcomes as well as the safety of the intervention.</p>

1.2.1.6. Vas 2009 ☆☆

Vas J et al. Correction of nonvertex presentation with moxibustion: a systematic review and metaanalysis. Am J Obstet Gynecol. 2009;201(3):241-59. [154014].

<p>Purpose</p>	<p>We searched systematically for randomized controlled trials, comparing moxibustion with a nonmoxibustion control group or other methods such as external cephalic version, postural methods, and acupuncture in databases, both Western and Chinese, up to June 2007.</p>
<p>Methods</p>	<p>Six studies, with 1087 subjects and a high degree of heterogeneity, compared moxibustion vs observation or postural methods. and reported a rate of cephalic version among the moxibustion group of 72.5% vs 53.2% in the control group (relative risk, 1.36; 95% confidence interval, 1.17-1.58); the number needed to treat was 5 (95% confidence interval, 4-7). In terms of safety, no significant differences were found in the comparison of moxibustion with other techniques.</p>
<p>Results</p>	<p>These studies reported a rate of cephalic version among the moxibustion group of 72.5% vs 53.2% in the control group (relative risk, 1.36; 95% confidence interval, 1.17-1.58); the number needed to treat was 5 (95% confidence interval, 4-7). In terms of safety, no significant differences were found in the comparison of moxibustion with other techniques.</p>
<p>Conclusion</p>	<p>Moxibustion at acupuncture point BL67 has been shown to produce a positive effect, whether used alone or in combination with acupuncture or postural measures, in- comparison with observation or postural methods alone, for the correction of nonvertex presentation, although these results should be viewed with caution, given the considerable. heterogeneity,found among-studies.</p>

1.2.1.7. Coyle 2005 ∅

Coyle M, Smith C, Peat B. Cephalic version by moxibustion for breech presentation. Cochrane

Database Syst Rev. 2005. (2):CD003928. [136184].

Objectives	To examine the effectiveness and safety of moxibustion on changing the presentation of an unborn baby in the breech position, the need for external cephalic version (ECV), mode of birth, and perinatal morbidity and mortality for breech presentation.
Methods	Search strategy: We searched the Cochrane Pregnancy and Childbirth Group trials register (30 August 2004), the Cochrane Central Register of Controlled Trials (The Cochrane Library, Issue 1, 2004), MEDLINE (1966 to March 2004), EMBASE (1980 to March 2004), CINAHL (1982 to March 2004), MIDIRS (1982 to March 2004), CISCOM (9 March 2004) and bibliographies of relevant papers. Selection CRITERIA: The inclusion criteria were published and unpublished randomised controlled trials comparing moxibustion (either alone or in combination with acupuncture) with a control group (no moxibustion), or other methods (e.g. external cephalic version, acupuncture) in women with a singleton breech presentation. Data collection and analysis: Both authors assessed eligibility and quality of trials independently. The outcome measures were baby's presentation at birth, need for external cephalic version, mode of birth, perinatal morbidity and mortality, maternal complications and maternal satisfaction, and adverse events.
Main results	Three trials involving a total of 597 women were included. Due to differences in interventions and sample size it was not appropriate to perform a meta-analysis for the main outcome. Only one trial reported on other outcome measures relevant to this review. Moxibustion reduced the need for ECV (relative risk (RR) 0.47, 95% confidence interval (CI) 0.33 to 0.66) and resulted in decreased use of oxytocin before or during labour for women who had vaginal deliveries (RR 0.28, 95% CI 0.13 to 0.60).
Authors' conclusions	There is insufficient evidence to support the use of moxibustion to correct a breech presentation. Moxibustion may be beneficial in reducing the need for ECV, and decreasing the use of oxytocin, however there is a need for well-designed randomised controlled trials to evaluate moxibustion for breech presentation which report on clinically relevant outcomes as well as the safety of the intervention.

2. Cost-effectiveness

See [corresponding item](#)

3. Clinical Practice Guidelines

⊕ positive recommendation (regardless of the level of evidence reported)
 ∅ negative recommendation, (or lack of evidence)

3.1. National Institute for Health and Care Excellence (NICE, UK) 2021 ∅

NICE guideline NG201 : Antenatal care [M] Management of breech presentation. National Institute for Health and Care Excellence (NICE). 2021. [218449]. [URL](#)

Complementary therapy: No evidence was identified. [Note: only 1 study included (acupuncture), all moxibustion studies excluded].

3.2. Danish Society of Obstetrics and Gynecology (DSOG, Denmark) 2020 ⊕

Breech delivery and external cephalic version. Danish Society of Obstetrics and Gynecology (DSOG).

2020;74P. [182690]. [URL](#)

There is limited evidence for the effect of Moxa, but a positive effect on fetal reversal from lower body to head position cannot be ruled out. (1a)

3.3. Collège national des gynécologues et obstétriciens français (CNGOF, France) 2019 Ø

Ducarme G. Présentation du siège. Recommandations pour la pratique clinique du CNGOF — Version par manœuvre externe et techniques de version alternatives. Gynecol Obstet Fertil Senol. 2020;48(1):81-94. [201194].

Sentilhes L, Schmitz T, Azria E, Gallot D, Ducarme G, Korb D, Mattuizzi A, Parant O, Sananès N, Baumann S, Rozenberg P, Sénat MV, Verspyck E. Présentation du siège . Recommandations pour la pratique clinique du CNGOF – Texte court Gynecol Obstet Fertil Senol. 2019. [200905].

En cas de présentation du siège, l'acupuncture, la moxibustion, les méthodes posturales (position genu-pectorale et pont indien) n'ont pas montré leur efficacité pour réduire le nombre de présentations du siège à la naissance (NP2), et ne sont pas recommandées (Grade B).

3.4. Royal College of Obstetricians and Gynaecologists (RCOG, UK) 2017 ☉

Impey LWM, Murphy DJ, Griffiths M, Penna LK on behalf of the Royal College of Obstetricians and Gynaecologists. External Cephalic Version and Reducing the Incidence of Term Breech Presentation. BJOG. 2017;124:e178-e192. [201232].

Women may wish to consider the use of moxibustion for breech presentation at 33–35 weeks of gestation, under the guidance of a trained practitioner.

4. Overviews of Clinical Practice Guidelines

4.1. Tsakiridis 2019

Tsakiridis I, Mamopoulos A, Athanasiadis A, Dagklis T. Management of Breech Presentation: A Comparison of Four National Evidence-Based Guidelines. Am J Perinatol. 2019. [201249].

OBJECTIVE: The management of breech presentation may improve perinatal outcomes. The aim of this study was to synthesize and compare published evidence of four national guidelines on breech presentation. **STUDY DESIGN:** A descriptive review of four recently published national guidelines on breech presentation and external cephalic version (ECV) was conducted: Royal College of Obstetricians and Gynaecologists guideline on “External Cephalic Version and Reducing the Incidence of Term Breech Presentation” and “Management of Breech Presentation”, American College of Obstetricians and Gynecologists guideline on “External Cephalic Version” and “Mode of Term Singleton Breech Delivery”. Society of Obstetricians and Gynaecologists of Canada guideline on “Vaginal Delivery of Breech Presentation” and the Royal Australian and New Zealand College of Obstetricians and Gynaecologists guideline on “Management of breech presentation at term.” **RESULTS:** Regarding ECV, there is no recommendation by the SOGC, whereas all other national guidelines recommend this technique. Regarding breech vaginal delivery, there are limited recommendations by the ACOG, whereas all other guidelines provide similar recommendations. The RANZCOG makes no special recommendations on the intrapartum period. **CONCLUSION:** The differences among national guidelines point out the need for the adoption of an international consensus on the management of breech presentation.

5. Randomized Controlled Trials

5.1. Sources

1. **Acudoc2**: base de données du Centre de Documentation du GERA (ECR non inclus dans les revues systématiques sources).
2. **Coyle 2023**: Coyle ME, Smith C, Peat B. Cephalic version by moxibustion for breech presentation. *Cochrane Database Syst Rev.* 2023 May 9;5:CD003928. <https://doi.org/10.1002/14651858.cd003928.pub4>
3. **Liao 2021** : Liao JA, Shao SC, Chang CT, Chai PYC, Owang KL, Huang TH, Yang CH, Lee TJ, Chen YC. Correction of Breech Presentation with Moxibustion and Acupuncture: A Systematic Review and Meta-Analysis. *Healthcare.* 2021;9(6):619. [212237]. <https://doi.org/10.3390/healthcare9060619>
4. **Zhang 2019**: Zhang Lu, Sun Weipeng, Cai Yinhe, Lu Ke, Yang Yaqin, Zeng Lei, Chen Baoyan. [Meta-analysis of Efficacy of Moxibustion at Zhiyin Point for Breech Presentation]. *Journal of Hubei University of Chinese Medicine.* 2019;5:117-21. [100163].
5. **Zhang 2013** : Zhang QH, Yue JH, Liu M, Sun ZR, Sun Q, Han C et al. Moxibustion for the correction of nonvertex presentation: a systematic review and meta-analysis of randomized controlled trials. *Evid Based Complement Alternat Med* 2013. [160352].
6. **Coyle 2012**: Coyle Me, Smith Ca, Peat B. Cephalic version by moxibustion for breech presentation. *Cochrane Database Syst Rev.* 2012. May 16:. [159289].
7. **Li 2009**: Li X, Hu J, Wang X, Zhang H, Liu J. Moxibustion and other acupuncture point stimulation methods to treat breech presentation: a systematic review of clinical trials. *Chin Med.* 2009;4(1):4.[153149].
8. **Liu 2009**: Liu Mai-Lan, Lan Lei, Tang Yong, Liang Fan-Rong. [Acupuncture and moxibustion for breech presentation: a systematic review]. *Chinese Journal of EBM.* 2009;9(8):840-3. [167341].
9. **Vas 2009**: Vas J et al. Correction of nonvertex presentation with moxibustion: a systematic review and metaanalysis. *Am J Obstet Gynecol.* 2009;201(3):241-59. [154014].
10. **Van Den Berg 2008**: Van Den Berg I et al. Effectiveness of acupuncture-type interventions versus expectant management to correct breech presentation: a systematic review. *Complement Ther Med.* 2008;16(2):92-100.149428
11. **Coyle 2005**: Coyle M, Smith C, Peat B. Cephalic version by moxibustion for breech presentation. *Cochrane Database Syst Rev.* 2005. (2):CD003928. [136184].

5.2. List

	Références	Comparisons	Sources
2021	Higashihara A, Horiuchi S. Smoke or smokeless moxibustion treatment for breech presentation: A three-arm pilot trial. <i>Jpn J Nurs Sci.</i> 2021 Oct;18(4):e12426. https://doi.org/10.1111/jjns.12426		Coyle 2023
2018	Qi Hong-yan. [Observations on the Efficacy of Point Zhiyin (BL67) Moxibustion plus Different Postures in Treating Fetal Malpositions]. <i>Shanghai Journal of Acupuncture and Moxibustion.</i> 2018;37(7):777. [182526].	Moxibustion zhiyin + posture A vs Moxibustion zhiyin + posture B vs Moxibustion zhiyin + posture C vs Moxibustion zhiyin + posture D	Acudoc2

	Références	Comparisons	Sources
2017	Keiko Tsujiuchi, Yoshihiko Koido, Syuuichi Katai, Hiromi Yoshikata. [Investigation of the effectiveness of acupuncture and moxibustion therapy as a treatment for breech presentation by a randomized controlled trial]. Journal of the Japan Society of Acupuncture and Moxibustion. 2017;67(3):215-223. [189633].		Coyle 2023
2016	Bue L, Lauszus FF. Moxibustion did not have an effect in a randomised clinical trial for version of breech position. Dan Med J. 2016;63(2). [189419].	Moxibustion zhiyin vs absence d'intervention	Coyle 2023, Liao 2021
	Cai Zhimin. [Clinical study of knee-thorax lying position combined with moxibustion to correct abnormal fetal position]. Asia Pacific Traditional Medicine. 2016;12(23):131-2. [99792].		Zhang 2019
	Sananes N, Roth GE, Aissi GA, Meyer N, Bigler A, Bouschbacher JM, Helmlinger C, Viville B, Guilpain M, Gaudineau A, Akladios CY, Nisand I, Langer B, Vayssiere C, Favre R. Acupuncture version of breech presentation: a randomized sham-controlled single-blinded trial. Eur J Obstet Gynecol Reprod Biol. 2016;;24-30. [187768].	Aiguille chauffée vs acupuncture factice	Acudoc2. Exclue de Coyle 2023 : The trial did not test moxibustion.
	Ye Liya, Wu Lei. [Outcome Analysis of Moxibustion Treatment of Buttock Pregnancy], For All Health (academic Edition) 大家健康 (学术版). 2016;10(6) : 42-3. [190575].		Zhang 2019
2015	Zhang Yaoxiu. [Analysis of the effects of moxibustion to Zhiyin point in correcting malposition in 63 cases]. Qinghai Medical Journal. 2015;45(12):53-4. [99960].	Moxibustion zhiyin vs posture	Zhang 2019
2014	Ai Xia, Gao Qiangqiang. [Correction Abnormal Fetal Position with Knee Chest and Moxibustion]. China Journal of Chinese Medicine. 2014;6:911-2. [111493].		Acudoc2
	Coulon C, Poleszczuk M, Paty-Montaigne Mh, Gascard C, Gay C, Houfflin-Debarge V, Subtil D. Version of breech fetuses by moxibustion with acupuncture: a randomized controlled trial. Obstet Gynecol. 2014;124(1):32-9. [177317].		Coyle 2023
2013	Vas J, Aranda-Regules JM, Modesto M, Ramos-Monserrat M, Barón M, Aguilar I, Benítez-Parejo N, Ramírez-Carmona C, Rivas-Ruiz F. Using moxibustion in primary healthcare to correct non-vertex presentation: a multicentre randomised controlled trial. Acupuncture in Medicine. 2013;31(1):31-8. [159287].	Moxibustion zhiyin vs moxibustion yinbai vs absence intervention	Coyle 2023, Liao 2021, Zhang 2019, Zhang 2013
	Yao Zuo , Li Chunxiang. [Correction of 52 cases of fetal malposition by moxibustion to Zhiyin point]. Zhejiang Journal of Traditional Chinese Medicine. 2013;48(11):828. [99992].	Moxibustion zhiyin vs absence d'intervention	Zhang 2019
2011	Do CK, Smith CA, Dahlen H, Bisits A, Schmied V. Moxibustion for cephalic version: a feasibility randomised controlled trial. BMC Complement Altern Med. 2011. [201312].	Moxibustion zhiyin vs absence d'intervention	Coyle 2023, Liao 2021, Zhang 2013

	Références	Comparisons	Sources
	Li Li. [Clinical Observation of Moxibustion to Zhiyin Point Combined with Knee-thoracic Lying Position Correction of Hip Position]. Chinese Community Physician: Medical Specialty. 2011;13(23):166. [100079].		Zhang 2019
2010	An Liqun. [Observation on 80 Cases of Correcting Pregnancy Buttocks with Moxibustion to Zhiyin Point Combined with Knee-Chest Position]. Zhejiang Journal of Traditional Chinese Medicine. 2010;45(2):139. [100699].		Zhang 2019
	Chen Ying, Yang Weijie, Cao Jingjing. [Clinical Observation on Treatment of Malpresentation by Mildness Moxibustion for 100 Cases]. Guangming Journal of Chinese Medicine. 2010;25(5):816-7. [111463].		Acudoc2
	Huo Ruixia. [Observation on effect of combined He-Ne laser irradiation with pure traditional Chinese herbs moxibustion of Zhiyin acupoint to correct parturient with breech presentation]. Chinese Nursing Research. 2010;22:2015-6. [111465].		Acudoc2
	Yang, Y.; Yin, S.C.; Syu, S.Y. [Integration Treatment of Traditional Chinese Medicine and Western Medicine in Breech Presentation: A Clinical Observation of 60 Cases]. J. Shanhai Med Coll. Contin. Educ. 2010, 20, 54-55		Liao 2021
2009	Guittier Mj, Pichon M, Dong H, Irion O, Boulvain M. Moxibustion for breech version: a randomized controlled trial. Obstet Gynecol. 2009;114(5):1034-40. [155295]. Guittier Mj, Klein Tj, Dong H, Andreoli N, Irion O, Boulvain M. Side-effects of moxibustion for cephalic version of breech presentation. Journal of Alternative and Complementary Medicine. 2008;14(10):1231-3.. [151207].	Moxibustion zhiyin vs absence d'intervention	Coyle 2023, Liao 2021, Zhang 2013, Coyle 2012
	Millereau M, Branger B, Darcel F. Étude comparative de la version foétale par acupuncture (moxibustion) versus groupe témoin. Journal de gynécologie, obstétrique et biologie de la reproduction.2009.38(6):481-487. [153316] .	Moxibustion zhiyin vs absence d'intervention	Coyle 2023, Liao 2021
2008	Yang Huilin, Wei Jianhong, Che Lifan, Zhang Yazuo, Cheng Meihong. [Application of Moxibustion to Zhiyin Point in Breech Presentation]. Chinese Journal for Clinicians. 2008;36(7):48-9. [100778].		Liao 2021, Zhang 2019
2007	Chen Y. [Moxibustion on zhiyin acupoint for treatment of 80 cases with fetal malposition]. Shaanxi Journal of TCM. 2007;28(3):334-5. [200856].	Moxibustion zhiyin vs posture	Liao 2021, Vas 2009
	Neri I, De Pace V, Venturini P, Facchinetti F. Effects of three different stimulations (acupuncture, moxibustion, acupuncture plus moxibustion) of bl.67 acupoint at small toe on fetal behavior of breech presentation. American Journal of Chinese Medicine. 2007;35(1):27-33. [145393]	Acupuncture vs Moxibustion vs acupuncture + moxibustion.	Coyle 2023, Coyle 2012, Vas 2009

	Références	Comparisons	Sources
	Wang Yan. [A Validation Study on Correcting Fetal Symptoms with Ai Tiao Moxibustion to Zhiyin Points]. Chengdu: Chengdu University of Traditional Chinese Medicine. 2007. [104372].		Zhang 2019
	Yang Yunkuan, Mao Min, Hu Youping, et al. [Effect of Moxibustion at Zhiyin (BL 67) to Correct the Fetus Malposition: Multi-center Randomized Controlled Clinical Study]. Journal of Traditional Chinese Medicine. 2007;48(12):1097-11000. [117976].	Moxibustion zhiyin vs posture	Zhang 2019, Zhang 2013, Liu 2009
2006	Li Yun. [Clinical observations on treatment of fetal malposition by suspended moxibustion over sanyinjiao]. Shanghai Journal of Acupuncture and Moxibustion. 2006;25(12):11. [145250].	Moxibustion zhiyin vs moxibustion sanyinjiao	Acudoc2
	Peng SM. [Comparison of several methods for conversion of breech presentation]. Zhongguo Yiyao Luntan. 2006;4:30. [201260].	Moxibustion zhiyin vs posture	Li 2009
	Yang Fengqin. [Knee chest position with moxibustion to the Yin point and simple knee chest position to correct the breech position]. Sichuan Traditional Chinese Medicine. 2006;24(5):106-7. [200852].	Moxibustion zhiyin + posture vs posture	Coyle 2023, Liao 2021, Zhang 2019, Zhang 2013, Coyle 2012, Li 2009, Liu 2009, Vas 2009
2005	Cardini F, Lombardo P, Regalia AI, Regaldo G, Zanini A, Negri Mg, Panepuccia L, Todros T. A randomised controlled trial of moxibustion for breech presentation. BJOG. 2005;112(6):743-7. [136456].	Moxibustion zhiyin vs absence d'intervention	Coyle 2023, Liao 2021, Zhang 2013, Coyle 2012, Li 2009, Liu 2009, Vas 2009 , Van den Berg 2008,
	Du Yanhua, Xue Aijing. [Clinical observation of 50 cases of correcting breech presentation with integrated traditional Chinese and western medicine]. Modern Journal of Integrated Traditional Chinese and Western Medicine. 2005;14(20):2727. [201267].	Moxibustion	Liu 2009
2004	Chen Y, Yang LW. [Moxibustion on Zhiyin plus raising buttocks in a lateral position for correction fetal presentation in 73 cases]. Zhongyiyao Linchuang Zazhi. 2004;16:333. [201306].	Moxibustion zhiyin + posture vs posture	Liao 2021, Coyle 2012, Li 2009
	Neri A, Airola G, Contu, Allais G, Facchinetti A. Acupuncture plus moxibustion to resolve breech presentation: a randomized controlled study. J Matern Fetal Neonatal Med. 2004;15:247-52. [129289].	Acupuncture + moxibustion zhiyin vs absence d'intervention	Coyle 2023, Liao 2021, Coyle 2012, Li 2009, Liu 2009, Vas 2009 , Van den Berg 2008, Coyle 2005
2003	Habek D, Cerkez Habek J, Jagust M. Acupuncture conversion of fetal breech presentation. Fetal Diagn Ther. 2003;18(6):418-21. [126110].	Acupuncture zhiyin vs absence intervention	Liao 2021, Li 2009, Liu 2009, Van den Berg 2008, Coyle 2005

Références		Comparisons	Sources
2002	Lin Yuanping, Zhang Deqing, Hao Yongqing, et al. [Combination of moxibustion at point zhiyin and knee-chest position for correction of breech pregnancy in 63 cases]. Chinese Acupuncture and Moxibustion. 2002;22(12):811. [110263].	Moxibustion zhiyin + posture vs posture	Liao 2021, Coyle 2012, Li 2009, Vas 2009
1998	Cardini F, Weixin H. Moxibustion for correction of breech presentation. JAMA.1998.280(18):1580-84. [58578].	Moxibustion zhiyin vs absence d'intervention	Coyle 2023, Liao 2021, Zhang 2013, Coyle 2012, Li 2009, Liu 2009, Vas 2009, Van den Berg 2008, Coyle 2005
	Ye LL, Zhu YP. [He-Ne laser stimulation for correcting breech presentaion in 133 cases]. Xiandai Fuchanke Jinzhan. 1998;7:379. [200814].	Laser zhiyin vs posture	Li 2009
1996	Li Qinghua et al. Clinical observation on correcting malposition of fetus by electro-acupuncture. Journal of TCM. 1996;16(4):260-2. [56846].	Electroacupuncture zhiyin vs moxibustion zhiyin vs absence d'intervention	Liao 2021, Liu 2009, Van den Berg 2008, Coyle 2005
1990	Huang WX. [Comparison study on moxibustion and knee-chest position for conversion of fetal presentation]. Zhongguo Zhongxiyi Jiehe Zazhi. 1990;10:105-6. [201268].	Moxibustion zhiyin vs posture vs absence d'intervention	Zhang 2019, Exclu Coyle 2012 ""as the method of group allocation was not reported, the decision was made to exclude the trial from the review"".

From:

<http://www.wiki-mtc.org/> - Encyclopédie des sciences médicales chinoises

Permanent link:

<http://www.wiki-mtc.org/doku.php?id=acupuncture:evaluation:gyneco-obstetrique:10.%20version%20foetale>



Last update: 04 May 2024 06:45