

Table des matières

1. Systematic Reviews and Meta-Analysis	1
1.1. Generic Acupuncture	1
1.1.1. Qin 2020	1
1.1.2. Beerepoot 2013 ☆	1
1.2. Special Clinical Forms	2
1.2.1. Recurrent Urinary Tract Infections in Pregnancy	2
2. Overviews of systematic reviews	2
2.1. Santer 2026	2
3. Clinical Practice Guidelines	3
3.1. Ministry of Public Health of Qatar (MOPH, Qatar) 2020 ⊕	3
3.2. Scottish Intercollegiate Guidelines Network (SIGN, Scotland) 2020 ⊕	3
3.3. Society of Obstetricians and Gynaecologists of Canada ((SOGC, Canada) 2017 ⊕	3
3.4. Société des Obstétriciens et Gynécologues du Canada (SOGC, Canada) 2010 ⊕	4
4. Randomized Controlled Trials	4
4.1. Sources	4
4.2. List	4

Recurrent urinary tract infections

Infections urinaires récidivantes : évaluation de l'acupuncture

1. Systematic Reviews and Meta-Analysis

1.1. Generic Acupuncture

1.1.1. Qin 2020

Qin X, Coyle ME, Yang L, Liang J, Wang K, Guo X, Zhang AL, Mao W, Lu C, Xue CC, Liu X. Acupuncture for recurrent urinary tract infection in women: A systematic review and meta-analysis. BJOG. 2020. [202654].

Background	Increasing antibiotic resistance has motivated interest in non-antibiotic prophylaxis of recurrent urinary tract infections (Ruti).
Objectives	To conduct a systematic review of the current state of evidence of acupuncture for uncomplicated Ruti in women.
Methods	SEARCH STRATEGY: Nine databases (PubMed, Embase, CENTRAL, CINAHL, AMED, CBM, CNKI, CQVIP, WANFANG) were searched from inception to February 2019. SELECTION CRITERIA: Randomised controlled trials (RCTs) evaluating the effects of acupuncture and related therapies for prophylaxis or treatment of uncomplicated Ruti in women were included. DATA COLLECTION AND ANALYSIS: Risk of bias was assessed, and the quality and strength of evidence evaluated using the GRADE framework. Results were reported as risk ratios (RR) for dichotomous outcomes or mean differences (MD) for continuous outcomes, with 95% confidence intervals (CI).
Main results	Five RCTs involving 341 participants were included. Methodological quality of studies and strength of the evidence were low to moderate. The chance of achieving composite cure with acupuncture therapies was greater than that with antibiotics (3 studies, 170 participants, RR 1.92, 95% CI 1.31 to 2.81, I ² =38%). The risk of UTI recurrence was lower with acupuncture compared with no treatment (2 studies, 135 participants, RR 0.39, 95% CI 0.26 to 0.58, I ² =0%) and sham acupuncture (1 study, 53 participants, RR 0.45, 95% CI 0.22 to 0.92).
Conclusions	Acupuncture appeared to be beneficial for treatment and prophylaxis of rUTIs, noting the limitations of the current evidence. Given the growing challenge of antibiotic resistance, there is a need for high-quality RCTs of non-pharmacological interventions such as acupuncture.

1.1.2. Beerepoot 2013 ☆

Beerepoot MA, Geerlings SE, Van Haarst EP, Van Charante NM, Ter Riet G. Nonantibiotic prophylaxis for recurrent urinary tract infections: a systematic review and meta-analysis of randomized controlled trials. J Urol. 2013. 90(6):1981-9. [170499].

Purpose	Increasing antimicrobial resistance has stimulated interest in nonantibiotic prophylaxis of recurrent urinary tract infections. We assessed the effectiveness, tolerability and safety of nonantibiotic prophylaxis in adults with recurrent urinary tract infections.
Materials and methods	MEDLINE®, EMBASE™, the Cochrane Library and reference lists of relevant reviews were searched to April 2013 for relevant English language citations. Two reviewers selected randomized controlled trials that met the predefined criteria for population, interventions and outcomes. The difference in the proportions of patients with at least 1 urinary tract infection was calculated for individual studies, and pooled risk ratios were calculated using random and fixed effects models. Adverse event rates were also extracted. The Jadad score was used to assess risk of bias (0 to 2-high risk and 3 to 5-low risk).
Results	We identified 5,413 records and included 17 studies with data for 2,165 patients. The oral immunostimulant OM-89 decreased the rate of urinary tract infection recurrence (4 trials, sample size 891, median Jadad score 3, RR 0.61, 95% CI 0.48-0.78) and had a good safety profile. The vaginal vaccine Urovac® slightly reduced urinary tract infection recurrence (3 trials, sample size 220, Jadad score 3, RR 0.81, 95% CI 0.68-0.96) and primary immunization followed by booster immunization increased the time to reinfection. Vaginal estrogens showed a trend toward preventing urinary tract infection recurrence (2 trials, sample size 201, Jadad score 2.5, RR 0.42, 95% CI 0.16-1.10) but vaginal irritation occurred in 6% to 20% of women. Cranberries decreased urinary tract infection recurrence (2 trials, sample size 250, Jadad score 4, RR 0.53, 95% CI 0.33-0.83) as did acupuncture (2 open label trials, sample size 165, Jadad score 2, RR 0.48, 95% CI 0.29-0.79) . Oral estrogens and lactobacilli prophylaxis did not decrease the rate of urinary tract infection recurrence.
Conclusions	The evidence of the effectiveness of the oral immunostimulant OM-89 is promising. Although sometimes statistically significant, pooled findings for the other interventions should be considered tentative until corroborated by more research. Large head-to-head trials should be performed to optimally inform clinical decision making.

1.2. Special Clinical Forms

1.2.1. Recurrent Urinary Tract Infections in Pregnancy

See [corresponding item](#)

2. Overviews of systematic reviews

2.1. Santer 2026

Santer M, Miller S, Fraser S, Francis N, Lown M, Becque T, Tenters F, Hanna L, Muller I. Non-antibiotic interventions to prevent recurrent urinary tract infections in women: overview of systematic reviews of randomised controlled trials. Br J Gen Pract. 2026 Jun 1:BJGP.2026.0055.

<https://doi.org/10.3399/BJGP.2026.0055>

Background	Recurrent urinary tract infections (rUTI), i.e. 2 or more UTIs in last 6 months or 3 or more in last 12 months, are common, have substantial impact on quality of life and cause major antibiotic use.
-------------------	--

Design	Overview of systematic reviews of randomised controlled trials (RCTs) of non-pharmaceutical and pharmaceutical (non-antibiotic) interventions for preventing rUTI in women.
Methods	We searched MEDLINE, EMBASE, CINAHL, Google Scholar and Cochrane from January 2014 to May 2025 for systematic reviews of RCTs assessing effectiveness of community-based non-antibiotic interventions for preventing UTI in women aged 16 or over with rUTI. We excluded studies in intensive care units, people with indwelling catheters or comparing different oral antibiotic regimens. There were no limitations regarding outcome measures or language.
Findings	Twenty-seven systematic reviews met inclusion criteria evaluating probiotics (9), D-mannose (7), vaginal oestrogen (7), cranberry (6), methenamine hippurate (3), herbal medicine (3), increased fluid intake (2), systemic oestrogen (4), acupuncture (2), vaccines (2). Eight examined combination strategies. Cranberry and methenamine hippurate use are supported by systematic reviews, as is vaginal oestrogen amongst postmenopausal women, although questions remain around dose, duration and treatment adherence. Reviews of probiotics, D-mannose, herbal medicine, acupuncture and vaccines concluded that stronger evidence is needed. Increased fluid intake benefits women with low intake, but further research is needed on optimal intake. There was little information on treatment adherence or acceptability for many interventions.
Conclusions	For effective interventions, questions remain about optimal use and for many stronger evidence is needed.

3. Clinical Practice Guidelines

⊕ positive recommendation (regardless of the level of evidence reported)
 ∅ negative recommendation (or lack of evidence)

3.1. Ministry of Public Health of Qatar (MOPH, Qatar) 2020 ⊕

The diagnosis & management of lower urinary tract infection in adult women. National Clinical Guidelines. Ministry of Public Health of Qatar (MOPH). 2020;:35P. [222747]. [URL](#)

Consider acupuncture as an alternative method to prevent recurrent UTIs in women who are unresponsive to or intolerant of antibiotic prophylaxis [L1, RGB].

3.2. Scottish Intercollegiate Guidelines Network (SIGN, Scotland) 2020 ⊕

Management of suspected bacterial lower urinary tract infection in adult women. Scottish Intercollegiate Guidelines Network (SIGN). 2020:68P. [210817]. [URL](#)

Acupuncture. Based on two small RCTs, a Canadian guideline recommended that acupuncture may be considered as an alternative intervention in the prevention of recurrent UTI in women who are unresponsive or intolerant to antibiotic prophylaxis. While acupuncture is not routinely used in Scotland for this indication its use could be considered by individual patients.

3.3. Society of Obstetricians and Gynaecologists of Canada ((SOGC, Canada) 2017 ⊕

Epp A, Larochelle A. Recurrent Urinary Tract Infection. J Obstet Gynaecol Can. 2017;39(10):e422-e431. [197220].

10. Acupuncture may be considered as an alternative in the prevention of recurrent urinary tract infections in women who are unresponsive to or intolerant of antibiotic prophylaxis. (I-B).

3.4. Société des Obstétriciens et Gynécologues du Canada (SOGC, Canada) 2010 ⊕

Epp A, Larochelle A. Recurrent Urinary Tract Infection. J Obstet Gynaecol Can. 2010;32(11):1082-90. [165533].

Acupuncture may be considered as an alternative in the prevention of recurrent urinary tract infections in women who are unresponsive to or intolerant of antibiotic prophylaxis. (I-B)

4. Randomized Controlled Trials


4.1. Sources

1. **Qin 2020:** Qin X, Coyle ME, Yang L, Liang J, Wang K, Guo X, Zhang AL, Mao W, Lu C, Xue CC, Liu X. Acupuncture for recurrent urinary tract infection in women: A systematic review and meta-analysis. BJOG. 2020. [202654].
2. **Beerepoot 2013:** Beerepoot MA, Geerlings SE, Van Haarst EP, Van Charante NM, Ter Riet G. Nonantibiotic prophylaxis for recurrent urinary tract infections: a systematic review and meta-analysis of randomized controlled trials. J Urol. 2013. 90(6):1981-9. [170499].

4.2. List

2018	Liu JL, Luo Q, Liu XH, Lin L. [Observation on the clinical effect of external treatment of strong renal moxibustion on recurrent urinary tract infection], China Modern Doctor. 2018;56(29):116-8. [202983].	Qin 2020
2013	Hong JY, Li F, Liang XQ, Hou Z. [Efficacy observation on female chronic pyelonephritis treated with abdominal cluster-needling therapy]. Chinese Acupuncture and Moxibustion. 2013;33(4):303-5. [162588].	Qin 2020
2010	Yu Siming, Guo Dandan. [Treatment of 30 cases of chronic urinary tract infection in adult women with moxibustion and antibiotics]. Shandong Journal of TCM. 2010;29(9):621-2. [160480].	Qin 2020
2002	Alraek T, Soedal LI, Fagerheim SU, Digranes A, Baerheim A. Acupuncture treatment in the prevention of uncomplicated recurrent lower urinary tract infections in adult women. Am J Public Health. 2002;92(10):1609-11. [107367]. Alraek T, Baerheim A. The effect of prophylactic acupuncture treatment in women with recurrent cystitis: Kidney patients fare better. Journal of Alternative and Complementary Medicine. 2003;9(5):651-8.. [117504].	Qin 2020, Beerepoot 2013
1998	Aune A, Alraek T, LiHua H, Baerheim A. Acupuncture in the prophylaxis of recurrent lower urinary tract infection in adult women. Scandinavian Journal of Primary Health Care. 1998;16(1):37-39. [58390].	Qin 2020, Beerepoot 2013

From:
<http://www.wiki-mtc.org/> - **Encyclopédie des sciences médicales chinoises**

Permanent link:
<http://www.wiki-mtc.org/doku.php?id=acupuncture:evaluation:uro-nephrologie:05.%20infections%20urinaires%20recidivantes> 

Last update: **02 Jun 2026 15:38**