

Table des matières

1. Systematic Reviews and Meta-Analysis	1
1.1. Li 2022 ☆	1
1.1.1. Xu 2016 Ø	2

Balance Function after Stroke

Troubles de l'équilibre post-AVC : Evaluation de l'acupuncture

Articles connexes: - évaluation du [taiji-qigong](#) -

1. Systematic Reviews and Meta-Analysis

1.1. Li 2022 ☆

Li B, Zhao Q, Du Y, Li X, Li Z, Meng X, Li C, Meng Z, Chen J, Liu C, Cao B, Chi S. Cerebral Blood Flow Velocity Modulation and Clinical Efficacy of Acupuncture for Posterior Circulation Infarction Vertigo: A Systematic Review and Meta-Analysis. *Evid Based Complement Alternat Med.* 2022 Jun 28;2022:3740856. <https://doi.org/10.1155/2022/3740856>

Background	Vertigo is a cardinal symptom of posterior circulation infarction (POCI). Acupuncture is demonstrated to have a beneficial effect on posterior circulation infarction vertigo (PCIV). However, the mechanism of acupuncture therapy is not clarified. This study aims to assess the cerebral blood flow velocity modulation and clinical efficacy of acupuncture for PCIV patients.
Methods	We conducted this systematic review for clinical randomized controlled trials (RCTs) regarding acupuncture on PCIV. The study duration was from September 2020 to September 2021. We searched the PubMed, EMBASE, Cochrane Library, Web of Science, Chinese Biomedical Literature Database (CBM), China National Knowledge Infrastructure (CNKI), Wanfang Database, and VIP. The publication date was set from inception to August 31, 2020. Based on the inclusion and exclusion criteria, two researchers independently screened literature and extracted data including basic study information, intervention details, outcome details, and adverse events. Outcome measures included the blood flow velocities of vertebrobasilar arteries and the Clinical Effective Rate of posterior circulation infarction vertigo. Pooled data were presented as standardized mean differences (SMDs) and relative risks (RR), with 95% confidence intervals (CIs). The meta-analysis was conducted using Review Manager software version 5.3.0.
Results	A total of 20 eligible RCTs (1541 participants) were included in this review, which compared acupuncture therapy (1 RCT) or acupuncture combined with pharmaceutical therapy (19 RCTs) to pharmaceutical therapy in patients with posterior circulation infarction vertigo. 7 studies assessed the blood flow velocities of the basilar artery examined by Transcranial Doppler (TCD), 8 studies assessed the bilateral vertebral arteries, and 13 studies evaluated the Clinical Effective Rate of posterior circulation infarction vertigo. Meta-analysis results showed that blood flow velocities of the basilar artery (SMD = 0.58, 95% CI = 0.40-0.76; P < 0.05), left vertebral artery (SMD = 0.48, 95% CI = 0.22-0.73; P < 0.05), and right vertebral artery (SMD = 0.44, 95% CI = 0.19-0.69; P < 0.05) were significantly higher in the acupuncture group compared with the control group. Clinical Effective Rate (RR = 1.22, 95% CI = 1.15-1.29; P = 0.792) was significantly better in the acupuncture group compared with the control group.

Conclusions	This study shows that acupuncture therapy is useful in improving the blood flow velocity of vertebrobasilar arteries and Clinical Effective Rate in patients with posterior circulation infarction vertigo. However, double-blind, sham-controlled trials with large sample sizes are required to support our conclusions.
--------------------	--

1.1.1. Xu 2016 Ø

Xu Hua,Tang Wei. [Effectiveness of Acupuncture and Rehabilitation Training on Balance Function after Stroke:A Systematic Review] Journal of Clinical Acupuncture and Moxibustion. 2016;32(11):56. [188602].

Objectives	To assess the effectiveness of acupuncture combined with rehabilitation training in the treatment of balance dysfunction after stroke.
Methods	Retrieve from CBMdisc, CNKI, v1P and Wanfang database. Collect clinically randomized controlled trials of comparison of acupuncture combined with rehabilitation Training and simple rehabilitation training, and assess the quality of the included studies by using RevMan5. 2 software Meta-analysis.
Results	There were totally 14 literatures, involving 1103 patients , The Berg Balance Scale [SMD=6.10,95% CI(5.39,6.82) P<0.00001],FMA function evaluation [SMD =1.68,95% C1 (1.47,1.91)P <0.00001], bathel index [SMD =5.90,95% CI(4.78,7.02)P<z0.00001] , Modified Bathel Index [SMD = 1.48,95% CI(1.05,1.91) ,P <0. 0001] ,FMA motor function evaluation [SMD = 1. 07,95% CI(0.87 , 1. 27) P <0. 0001] , neural function defect score [SMD = - 1. 60 , 95% CI (- 2. 10, - 1. i 1) P < 0.0001]and the Barthel index were as the basis to evaluate the curative effect□RR:1.16,95% CI(0.98, 1. 37)P =0. 08□□There was no significant difference between two groups.
Conclusions	Acupuncture combined with rehabilitation training is effective in the treatment of balance disorder after stroke. Because the quality of the included studies is not high the conclusion still needs to be randomized controlled trials with high quality of strict design and large samples.

From:
<https://www.wiki-mtc.org/> - Encyclopédie des sciences médicales chinoises

Permanent link:
<https://www.wiki-mtc.org/doku.php?id=acupuncture:evaluation:neuro-psychiatrie:07.%20troubles%20de%20l%20equilibre%20post-avc>

Last update: **01 Dec 2025 15:10**